



Scientific Inquiry and Review (SIR)

Volume 4, Issue 4, December 2020

ISSN (P): 2521-2427, ISSN (E): 2521-2435

Journal DOI: <https://doi.org/10.32350/sir>

Issue DOI: <https://doi.org/10.32350/sir.44>

Homepage: <https://journals.umt.edu.pk/index.php/SIR/Home>

Journal QR Code:



Article **Zagreb Polynomials and Redefined Zagreb Indices for Chemical Structures Helpful in the Treatment of COVID-19**

Author(s)

Abaid ur Rehman Virk


Online
Published

December 2020

Article DOI

<https://doi.org/10.32350/sir.44.04>

QR Code
of Article



Abaid ur Rehman

To cite this
Article

Virk Abaid ur Rehman. Zagreb polynomials and redefined zagreb indices for chemical structures helpful in the treatment of COVID-19. *Sci Inquiry Rev.* 2020;4(4):46–62.

[Crossref](#)

Copyright
Information

This article is open access and is distributed under the terms of Creative Commons Attribution – Share Alike 4.0 International License.



A publication of the
School of Science, University of Management and Technology
Lahore, Pakistan.

Indexing
&
Abstracting



Zagreb Polynomials and Redefined Zagreb Indices for Chemical Structures Helpful in the Treatment of COVID-19

Abaid ur Rehman Virk

Department of Mathematics,
University of Sialkot, Sialkot, Pakistan

*abaid.math@gmail.com

Abstract

A topological index (TI) is a number that is helpful in predicting the properties of chemical compounds. We can estimate the physical and chemical properties of several chemical compounds. In this study, we compute Zagreb polynomials and the redefined Zagreb indices for chemical compounds used in the treatment of COVID-19 namely remdesivir, chloroquine, hydroxychloroquine and theaflavin.

Keywords: COVID-19, chloroquine, hydroxychloroquine, redefined Zagreb indices, remdesivir, theaflavin, Zagreb polynomials

Introduction

Currently, the COVID-19 pandemic is jeopardizing human health as well as the economy all over the world. Wuhan, China is its place of origin [1]. By December 2020, there were 80410714 confirmed cases of COVID including 1732088 deaths worldwide (as per world meter information). Scientists have used some existing antiviral agents to treat the disease [2, 3, 4, 5, 6]. These include chloroquine, hydroxychloroquine, remdesivir (GS5734), and theaflavin [7].

Topological Indices (TIs) are very important tools in predicting the various properties of chemical compounds. A special number in graph theoretical term which represents a molecular structure is known as a TI. The first and second Zagreb indices are the oldest molecular descriptors introduced by Gutman in 1975 [8] and their various properties are investigated in this research.

They are defined as follows:

$$M_1(G) = \sum_{uv \in E(G)} (d_u + d_v)$$

$$M_2(G) = \sum_{uv \in E(G)} (d_u \times d_v)$$

Considering the Zagreb indices in [9], Fath-Tabar introduced the first and second Zagreb polynomials which are stated as follows:

$$M_1(G, z) = \sum_{uv \in E(G)} z^{[d_u + d_v]}$$

$$M_2(G, z) = \sum_{uv \in E(G)} z^{[d_u \times d_v]}$$

$M_1(G, x)$ and $M_2(G, x)$ give us important information about molecular graphs [10]. Afterwards, the author in [11] defined the third Zagreb polynomial as follows:

$$M_3(G, z) = \sum_{uv \in E(G)} z^{[d_u - d_v]}$$

Other versions of the Zagreb polynomials are given in [12].

$$M_4(G, z) = \sum_{uv \in E(G)} z^{d_u [d_u + d_v]}$$

$$M_5(G, z) = \sum_{uv \in E(G)} z^{d_v [d_u + d_v]}$$

$$M_{a,b}(G, z) = \sum_{uv \in E(G)} z^{[ad_u + bd_v]}$$

$$M'_{a,b}(G, z) = \sum_{uv \in E(G)} z^{(d_u + a)(b + d_v)}$$

Ranjini et al. [13] introduced the redefined Zagreb indices as follows:

$$ReZG_1 = \sum_{uv \in E(G)} \frac{d_u + d_v}{d_u \times d_v}$$

$$ReZG_2 = \sum_{uv \in E(G)} \frac{d_u \times d_v}{d_u + d_v}$$

$$ReZG_3 = \sum_{uv \in E(G)} (d_u + d_v) (d_u \times d_v)$$

Many studies have investigated the topological invariants [14, 15, 16, 17, 18, 19].

2. Main Results

In this section, we will discuss Zagreb polynomials and the redefined Zagreb indices for chemical structures used in the treatment of COVID-19 namely remdesivir, chloroquine, hydroxychloroquine and theaflavin.

2.1. Zagreb Polynomials and Redefined Zagreb Indices for Remdesivir

The graph of remdesivir is given in Figure 1. There are eight types of edges present in the graph. The degree-based edge partition is given in Table 1.




Figure 1. Chemical structure of Remdesivir (National centre for biotechnology information (PubChem))

Table 1. Partition of E (Remdesivir)

(d_u, d_v)	Frequency
(1,2)	2
(1,3)	5
(1,4)	2
(2,2)	9
(2,3)	14
(2,4)	4
(3,3)	6
(3,4)	2

Theorem 2.2.

Let G be the graph of Remdesivir. The Zagreb polynomials are stated as follows:

- (1) $M_1(G, z) = 2z^3 + 14z^4 + 16z^5 + 10z^6 + 2z^7$
- (2) $M_2(G, z) = 2z^2 + 5z^3 + 11z^4 + 14z^6 + 4z^8 + 6z^9 + 2z^{12}$
- (3) $M_3(G, z) = 2z^3 + 9z^2 + 18z + 15$
- (4) $M_4(G, z) = 2z^3 + 5z^4 + 2z^5 + 9z^8 + 14z^{10} + 4z^{12} + 6z^{18} + 2z^{36}$
- (5) $M_5(G, z) = 2z^6 + 5z^{12} + 2z^{20} + 9z^8 + 14z^{15} + 4z^{24} + 6z^{18} + 2z^{48}$
- (6) $M_{a,b}(G, z) = 2z^{a+2b} + 5z^{a+3b} + 2z^{a+3b} + 9z^{2a+2b} + 14z^{2a+3b} + 4z^{2a+4b} + 6z^{3a+3b} + 2z^{3a+4b}$
- (7) $M'_{a,b}(G, z) = 2z^{(1+a)(2+b)} + 5z^{(1+a)(3+b)} + 2z^{(1+a)(4+b)} + 9z^{(2+a)(2+b)} + 14z^{(2+a)(3+b)} + 4z^{(2+a)(4+b)} + 6z^{(3+a)(3+b)} + 2z^{(3+a)(4+b)}$

Proofs:

1. $M_1(G, z) = \sum_{uv \in E(G)} z^{[d_u + d_v]}$
 $= |E_{\{1,2\}}(G)| z^3 + |E_{\{1,3\}}(G)| z^4 + |E_{\{1,4\}}(G)| z^5 + |E_{\{2,2\}}(G)| z^4 + |E_{\{2,3\}}(G)| z^5 +$
 $|E_{\{2,4\}}(G)| z^6 + |E_{\{3,3\}}(G)| z^6 + |E_{\{3,4\}}(G)| z^7$
 $= 2z^3 + 14z^4 + 16z^5 + 10z^6 + 2z^7$
2. $M_2(G, z) = \sum_{uv \in E(G)} z^{[d_u \times d_v]}$
 $= |E_{\{1,2\}}(G)| z^2 + |E_{\{1,3\}}(G)| z^3 + |E_{\{1,4\}}(G)| z^4 + |E_{\{2,2\}}(G)| z^4 + |E_{\{2,3\}}(G)| z^6 +$
 $|E_{\{2,4\}}(G)| z^8 + |E_{\{3,3\}}(G)| z^9 + |E_{\{3,4\}}(G)| z^{12}$
 $= 2z^2 + 5z^3 + 11z^4 + 14z^6 + 4z^8 + 6z^9 + 2z^{12}$
3. $M_3(G, z) = \sum_{uv \in E(G)} z^{[d_u - d_v]}$
 $= |E_{\{1,2\}}(G)| z^1 + |E_{\{1,3\}}(G)| z^2 + |E_{\{1,4\}}(G)| z^3 + |E_{\{2,2\}}(G)| z^0 + |E_{\{2,3\}}(G)| z^1 +$
 $|E_{\{2,4\}}(G)| z^2 + |E_{\{3,3\}}(G)| z^0 + |E_{\{3,4\}}(G)| z^1$
 $= 2z^3 + 9z^2 + 18z + 15$

$$\begin{aligned}
4. \quad M_4(G, z) &= \sum_{uv \in E(G)} z^{d_u[d_u+d_v]} \\
&= |E_{\{1,2\}}(G)| z^{3+|E_{\{1,3\}}(G)|} z^{4+|E_{\{1,4\}}(G)|} \\
&\quad z^{5+|E_{\{2,2\}}(G)|} z^{8+|E_{\{2,3\}}(G)|} z^{10+|E_{\{2,4\}}(G)|} z^{12+|E_{\{3,3\}}(G)|} z^{18+|E_{\{3,4\}}(G)|} z^{36} \\
&= 2z^3 + 5z^4 + 2z^5 + 9z^8 + 14z^{10} + 4z^{12} + 6z^{18} + 2z^{36} \\
5. \quad M_5(G, z) &= \sum_{uv \in E(G)} z^{d_v[d_u+d_v]} \\
&= |E_{\{1,2\}}(G)| z^{6+|E_{\{1,3\}}(G)|} z^{12+|E_{\{1,4\}}(G)|} \\
&\quad z^{20+|E_{\{2,2\}}(G)|} z^{8+|E_{\{2,3\}}(G)|} z^{15+|E_{\{2,4\}}(G)|} z^{24+|E_{\{3,3\}}(G)|} z^{18+|E_{\{3,4\}}(G)|} z^{48} \\
&= 2z^6 + 5z^{12} + 2z^{20} + 9z^8 + 14z^{15} + 4z^{24} + 6z^{18} + 2z^{48} \\
6. \quad M_{a,b}(G, z) &= \sum_{uv \in E(G)} z^{[ad_u+bd_v]} \\
&= |E_{\{1,2\}}(G)| z^{a+2b+|E_{\{1,3\}}(G)|} z^{a+3b+|E_{\{1,4\}}(G)|} \\
&\quad z^{a+4b+|E_{\{2,2\}}(G)|} z^{2a+2b+|E_{\{2,3\}}(G)|} z^{2a+3b+|E_{\{2,4\}}(G)|} z^{2a+4b+|E_{\{3,3\}}(G)|} z^{3a+3b+|E_{\{3,4\}}(G)|} z^{3a+4b} \\
&= 2z^{a+2b} + 5z^{a+3b} + 2z^{a+4b} + 9z^{2a+2b} + 14z^{2a+3b} + 4z^{2a+4b} + 6z^{3a+3b} + 2z^{3a+4b} \\
7. \quad M'_{a,b}(G, z) &= \sum_{uv \in E(G)} z^{(d_u+a)(b+d_v)} \\
&= |E_{\{1,2\}}(G)| z^{(1+a)(2+b)+|E_{\{1,3\}}(G)|} z^{(1+a)(3+b)+|E_{\{1,4\}}(G)|} z^{(1+a)(4+b)+|E_{\{2,2\}}(G)|} z^{(2+a)(2+b)+|E_{\{2,3\}}(G)|} z^{(2+a)(3+b)+|E_{\{2,4\}}(G)|} z^{(2+a)(4+b)+|E_{\{3,3\}}(G)|} z^{(3+a)(3+b)+|E_{\{3,4\}}(G)|} z^{(3+a)(4+b)} \\
&= 2z^{(1+a)(2+b)} + 5z^{(1+a)(3+b)} + 2z^{(1+a)(4+b)} + 9z^{(2+a)(2+b)} + 14z^{(2+a)(3+b)} + 4z^{(2+a)(4+b)} + 6z^{(3+a)(3+b)} + 2z^{(3+a)(4+b)}
\end{aligned}$$

Theorem 2.3.

Let G be the graph of Remdesivir. The redefined Zagreb indices are given below.

- (1) $ReZG_1 = 41$
- (2) $ReZG_2 = 52.64$
- (3) $ReZG_3 = 1948$

Proofs:

$$1. \text{ReZG}_1 = \sum_{uv \in E(G)} \frac{d_u + d_v}{d_u \times d_v}$$

$$= (2) \left(\frac{3}{2}\right) + (5) \left(\frac{4}{3}\right) + (2) \left(\frac{5}{4}\right) + (9) \left(\frac{4}{4}\right) + (14) \left(\frac{5}{6}\right) + (4) \left(\frac{6}{8}\right) + (6) \left(\frac{6}{9}\right) + (2) \left(\frac{7}{12}\right)$$

$$= 41$$

$$2. \text{ReZG}_2 = \sum_{uv \in E(G)} \frac{d_u \times d_v}{d_u + d_v}$$

$$= (2) \left(\frac{2}{3}\right) + (5) \left(\frac{3}{4}\right) + (2) \left(\frac{4}{5}\right) + (9) \left(\frac{4}{4}\right) + (14) \left(\frac{6}{5}\right) + (4) \left(\frac{8}{6}\right) + (6) \left(\frac{9}{6}\right) + (2) \left(\frac{12}{7}\right)$$

$$= 52.64$$

$$3. \text{ReZG}_3 = \sum_{uv \in E(G)} (d_u + d_v) (d_u \times d_v)$$

$$= (2) (2 \times 3) + (5) (3 \times 4) + (2) (4 \times 5) + (9) (4 \times 4) +$$

$$(14) (6 \times 5) + (4) (8 \times 6) + (6) (9 \times 6) + (9) (12 \times 7)$$

$$= 1948$$

2.4. Zagreb Polynomials and Redefined Zagreb Indices for Chloroquine

The graph of chloroquine is given in Figure 2. There are five types of edges present in the graph. The degree-based edge partition is given in Table 2.




Figure 2. Chemical structure of of Chloroquine (National centre for biotechnology information (PubChem))

Table 2. Partition of E (Chloroquine)

(du, dv)	Frequency
(1, 2)	2
(1,3)	2
(2,2)	5
(2,3)	12
(3,3)	2

Theorem 2.5.

Let G be the graph of Chloroquine. The Zagreb polynomials are stated as follows:

- (1) $M_1(G, z) = 2z^3 + 7z^4 + 12z^5 + 2z^6$
- (2) $M_2(G, z) = 2z^2 + 2z^3 + 5z^4 + 12z^6 + 2z^9$
- (3) $M_3(G, z) = 2z^2 + 14z + 7$
- (4) $M_4(G, z) = 2z^3 + 2z^4 + 5z^8 + 12z^{10} + 2z^{18}$
- (5) $M_5(G, z) = 2z^6 + 2z^{12} + 5z^8 + 12z^{15} + 2z^{18}$
- (6) $M_{a,b}(G, z) = 2z^{a+2b} + 2z^{a+3b} + 5z^{2a+2b} + 12z^{2a+3b} + 2z^{3a+3b}$
- (7) $M'_{a,b}(G, z) = 2z^{(1+a)(2+b)} + 2z^{(1+a)(3+b)} + 5z^{(2+a)(2+b)} + 12z^{(2+a)(3+b)} + 2z^{(3+a)(3+b)}$

Proofs:

$$\begin{aligned}
 M_1(G, z) &= \sum_{uv \in E(G)} z^{[d_u + d_v]} \\
 &= |E_{\{1,2\}}(G)| z^3 + |E_{\{1,3\}}(G)| z^4 + |E_{\{2,2\}}(G)| z^4 + |E_{\{2,3\}}(G)| z^5 + |E_{\{3,3\}}(G)| z^6 \\
 &= 2z^3 + 7z^4 + 12z^5 + 2z^6
 \end{aligned}$$

$$\begin{aligned}
 1. \quad M_2(G, z) &= \sum_{uv \in E(G)} z^{[d_u \times d_v]} \\
 &= |E_{\{1,2\}}(G)| z^2 + |E_{\{1,3\}}(G)| z^3 + |E_{\{2,2\}}(G)| z^4 + |E_{\{2,3\}}(G)| z^6 + |E_{\{3,3\}}(G)| z^9 \\
 &= 2z^2 + 2z^3 + 5z^4 + 12z^6 + 2z^9
 \end{aligned}$$

$$2. \quad M_3(G, z) = \sum_{uv \in E(G)} z^{[d_u - d_v]}$$

$$\begin{aligned}
&= |E_{\{1,2\}}(G)| z^1 + |E_{\{1,3\}}(G)| z^2 + |E_{\{2,2\}}(G)| \\
& z^0 + |E_{\{2,3\}}(G)| z^1 + |E_{\{3,3\}}(G)| z^0 \\
&= 2z^2 + 14z + 7
\end{aligned}$$

$$\begin{aligned}
3. \quad M_4(G, z) &= \sum_{uv \in E(G)} z^{d_u[d_u+d_v]} \\
&= |E_{\{1,2\}}(G)| z^3 + |E_{\{1,3\}}(G)| z^4 + |E_{\{2,2\}}(G)| \\
& z^8 + |E_{\{2,3\}}(G)| z^{10} + |E_{\{3,3\}}(G)| z^{18} \\
&= 2z^3 + 2z^4 + 5z^8 + 12z^{10} + 2z^{18}
\end{aligned}$$

$$\begin{aligned}
4. \quad M_5(G, z) &= \sum_{uv \in E(G)} z^{d_v[d_u+d_v]} \\
&= |E_{\{1,2\}}(G)| z^6 + |E_{\{1,3\}}(G)| z^{12} + |E_{\{2,2\}}(G)| \\
& z^8 + |E_{\{2,3\}}(G)| z^{15} + |E_{\{3,3\}}(G)| z^{18} \\
&= 2z^6 + 5z^8 + 2z^{12} + 12z^{15} + 2z^{18}
\end{aligned}$$

$$\begin{aligned}
5. \quad M_{a,b}(G, z) &= \sum_{uv \in E(G)} z^{[ad_u+bd_v]} \\
&= |E_{\{1,2\}}(G)| z^{a+2b} + |E_{\{1,3\}}(G)| \\
& z^{a+3b} + |E_{\{2,2\}}(G)| z^{2a+2b} + |E_{\{2,3\}}(G)| z^{2a+3b} \\
& + |E_{\{3,3\}}(G)| z^{3a+3b} \\
&= 2z^{a+2b} + 2z^{a+3b} + 5z^{2a+2b} + 12z^{2a+3b} + 2z^{3a+3b}
\end{aligned}$$

$$\begin{aligned}
6. \quad M'_{a,b}(G, z) &= \sum_{uv \in E(G)} z^{(d_u+a)(b+d_v)} \\
&= |E_{\{1,2\}}(G)| z^{(1+a)(2+b)} + |E_{\{1,3\}}(G)| z^{(1+a)(3+b)} + \\
& |E_{\{2,2\}}(G)| z^{(2+a)(2+b)} + |E_{\{2,3\}}(G)| z^{(2+a)(3+b)} + \\
& |E_{\{3,3\}}(G)| z^{(3+a)(3+b)} \\
&= 2z^{(1+a)(2+b)} + 2z^{(1+a)(3+b)} + 5z^{(2+a)(2+b)} + 12z^{(2+a)(3+b)} + 2z^{(3+a)(3+b)}
\end{aligned}$$

Theorem 2.6.

Let G be the graph of chloroquine. The redefined Zagreb indices are given below.

- (1) $ReZG_1 = 22$
- (2) $ReZG_2 = 25.23$
- (3) $ReZG_3 = 584$

Proofs:

$$1. \text{ReZG}_1 = \sum_{uv \in E(G)} \frac{d_u + d_v}{d_u \times d_v}$$

$$= (2) \left(\frac{3}{2}\right) + (2) \left(\frac{4}{3}\right) + (5) \left(\frac{4}{4}\right) + (12) \left(\frac{5}{6}\right) + (2) \left(\frac{6}{9}\right) = 22$$

$$2. \text{ReZG}_2 = \sum_{uv \in E(G)} \frac{d_u \times d_v}{d_u + d_v}$$

$$= (2) \left(\frac{2}{3}\right) + (2) \left(\frac{3}{4}\right) + (5) \left(\frac{4}{4}\right) + (12) \left(\frac{6}{5}\right) + (2) \left(\frac{9}{6}\right) = 25.23$$

$$3. \text{ReZG}_3 = \sum_{uv \in E(G)} (d_u + d_v) (d_u \times d_v)$$

$$= (2) (2 \times 3) + (5) (3 \times 4) + (9) (4 \times 4) + (14) (6 \times 5) + (6) (9 \times 6) = 584$$

2.7. Zagreb Polynomials and Redefined Zagreb Indices for Hydroxychloroquine.

The graph of hydroxychloroquine is given in Figure 3. There are five types of edges present in the graph. The degree-based edge partition is given in Table 3.




Figure 3. Chemical structure of Hydroxychloroquine (National centre for biotechnology information (PubChem))

Table 3. Partition of E (Hydroxychloroquine)

(du, dv)	Frequency
(1, 2)	2
(1,3)	2
(2,2)	6
(2,3)	12
(3,3)	2

Theorem 2.8.

Let G be the graph of hydroxychloroquine. The Zagreb polynomials are stated as follows:

- (1) $M_1(G, z) = 2z^3 + 8z^4 + 12z^5 + 2z^6$
- (2) $M_2(G, z) = 2z^2 + 2z^3 + 6z^4 + 12z^6 + 2z^9$
- (3) $M_3(G, z) = 2z^2 + 14z + 8$
- (4) $M_4(G, z) = 2z^3 + 2z^4 + 6z^8 + 12z^{10} + 2z^{18}$
- (5) $M_5(G, z) = 2z^6 + 6z^8 + 2z^{12} + 12z^{15} + 2z^{18}$
- (6) $M_{a,b}(G, z) = 2z^{a+2b} + 2z^{a+3b} + 6z^{2a+2b} + 12z^{2a+3b} + 2z^{3a+3b}$
- (7) $M'_{a,b}(G, z) = 2z^{(1+a)(2+b)} + 2z^{(1+a)(3+b)} + 6z^{(2+a)(2+b)} + 12z^{(2+a)(3+b)} + 2z^{(3+a)(3+b)}$

Proofs:

1.
$$\begin{aligned}
 M_1(G, z) &= \sum_{uv \in E(G)} z^{[d_u + d_v]} \\
 &= |E_{\{1,2\}}(G)| z^3 + |E_{\{1,3\}}(G)| z^4 + |E_{\{2,2\}}(G)| z^4 + |E_{\{2,3\}}(G)| z^5 + |E_{\{3,3\}}(G)| z^6 \\
 &= 2z^3 + 8z^4 + 12z^5 + 2z^6
 \end{aligned}$$
2.
$$\begin{aligned}
 M_2(G, z) &= \sum_{uv \in E(G)} z^{[d_u \times d_v]} \\
 &= |E_{\{1,2\}}(G)| z^2 + |E_{\{1,3\}}(G)| z^3 + |E_{\{2,2\}}(G)| z^4 + |E_{\{2,3\}}(G)| z^6 + |E_{\{3,3\}}(G)| z^9 \\
 &= 2z^2 + 2z^3 + 6z^4 + 12z^6 + 2z^9
 \end{aligned}$$
3.
$$M_3(G, z) = \sum_{uv \in E(G)} z^{[d_u - d_v]}$$

$$\begin{aligned}
&= |E_{\{1,2\}}(G)| z^1 + |E_{\{1,3\}}(G)| z^2 + |E_{\{2,2\}}(G)| \\
& z^0 + |E_{\{2,3\}}(G)| z^1 + |E_{\{3,3\}}(G)| z^0 \\
&= 2z^2 + 14z + 8
\end{aligned}$$

$$\begin{aligned}
4. \quad M_4(G, z) &= \sum_{uv \in E(G)} z^{d_u[d_u+d_v]} \\
&= |E_{\{1,2\}}(G)| z^3 + |E_{\{1,3\}}(G)| z^4 + |E_{\{2,2\}}(G)| \\
& z^8 + |E_{\{2,3\}}(G)| z^{10} + |E_{\{3,3\}}(G)| z^{18} \\
&= 2z^3 + 2z^4 + 6z^8 + 12z^{10} + 2z^{18}
\end{aligned}$$

$$\begin{aligned}
5. \quad M_5(G, z) &= \sum_{uv \in E(G)} z^{d_v[d_u+d_v]} \\
&= |E_{\{1,2\}}(G)| z^6 + |E_{\{1,3\}}(G)| z^{12} + |E_{\{2,2\}}(G)| \\
& z^8 + |E_{\{2,3\}}(G)| z^{15} + |E_{\{3,3\}}(G)| z^{18} \\
&= 2z^6 + 6z^8 + 2z^{12} + 12z^{15} + 2z^{18}
\end{aligned}$$

$$\begin{aligned}
6. \quad M_{a,b}(G, z) &= \sum_{uv \in E(G)} z^{[ad_u+bd_v]} \\
&= |E_{\{1,2\}}(G)| z^{a+2b} + |E_{\{1,3\}}(G)| \\
& z^{a+3b} + |E_{\{2,2\}}(G)| z^{2a+2b} + |E_{\{2,3\}}(G)| z^{2a+3b} \\
& + |E_{\{3,3\}}(G)| z^{3a+3b} \\
&= 2z^{a+2b} + 2z^{a+3b} + 6z^{2a+2b} + 12z^{2a+3b} + 2z^{3a+3b}
\end{aligned}$$

$$\begin{aligned}
7. \quad M'_{a,b}(G, z) &= \sum_{uv \in E(G)} z^{(d_u+a)(b+d_v)} \\
&= |E_{\{1,2\}}(G)| z^{(1+a)(2+b)} + |E_{\{1,3\}}(G)| z^{(1+a)(3+b)} + \\
& |E_{\{2,2\}}(G)| z^{(2+a)(2+b)} + |E_{\{2,3\}}(G)| z^{(2+a)(3+b)} + \\
& |E_{\{3,3\}}(G)| z^{(3+a)(3+b)} \\
&= 2z^{(1+a)(2+b)} + 2z^{(1+a)(3+b)} + 6z^{(2+a)(2+b)} + 12z^{(2+a)(3+b)} + 2z^{(3+a)(3+b)}
\end{aligned}$$

Theorem 2.9.

Let G be the graph of hydroxychloroquine. The redefined Zagreb indices are given below.

$$(1) \text{Re}ZG_1 = 23$$

$$(2) \text{Re}ZG_2 = 26.23$$

$$(3) \text{Re}ZG_3 = 600$$

Proofs:

$$\begin{aligned}
 1. \quad ReZG_1 &= \sum_{uv \in E(G)} \frac{d_u + d_v}{d_u \times d_v} \\
 &= (2) \left(\frac{3}{2}\right) + (2) \left(\frac{4}{3}\right) + (6) \left(\frac{4}{4}\right) + (12) \left(\frac{5}{6}\right) + (2) \left(\frac{6}{9}\right) \\
 &= 23
 \end{aligned}$$

$$\begin{aligned}
 2. \quad ReZG_2 &= \sum_{uv \in E(G)} \frac{d_u \times d_v}{d_u + d_v} \\
 &= (2) \left(\frac{2}{3}\right) + (2) \left(\frac{3}{4}\right) + (6) \left(\frac{4}{4}\right) + (12) \left(\frac{6}{5}\right) + (2) \left(\frac{9}{6}\right) \\
 &= 26.23
 \end{aligned}$$

$$\begin{aligned}
 3. \quad ReZG_3 &= \sum_{uv \in E(G)} (d_u + d_v) (d_u \times d_v) \\
 &= (2) (2 \times 3) + (2) (3 \times 4) + (6) (4 \times 4) + (12) (6 \times 5) + \\
 &\quad (2) (9 \times 6) = 600
 \end{aligned}$$

2.10. Zagreb Polynomials and Redefined Zagreb Indices for Theaflavin

The graph of theaflavin is given in Figure 4. There are three types of edges present in the graph. The degree-based edge partition is given in Table 4.




Figure 4. Chemical Structure of Theaflavin (National Centre for Biotechnology information (PubChem))

Table 4. Partition of E(Theaflavin)

(du,dv)	Frequency
(1,3)	10
(2,3)	22
(3,3)	14

Theorem 2.11.

Let G be the graph of theaflavin. The Zagreb polynomials are stated as follows:

- (1) $M_1(G, z) = 10z^4 + 22z^5 + 14z^6$
- (2) $M_2(G, z) = 10z^3 + 22z^6 + 14z^9$
- (3) $M_3(G, z) = 10z^2 + 22z + 14$
- (4) $M_4(G, z) = 10z^4 + 22z^{12} + 14z^{18}$
- (5) $M_5(G, z) = 10z^{12} + 22z^{15} + 14z^{18}$
- (6) $M_{a,b}(G, z) = 10z^{a+3b} + 22z^{2a+3b} + 14z^{3a+3b}$
- (7) $M'_{a,b}(G, z) = 10z^{(1+a)(3+b)} + 22z^{(2+a)(3+b)} + 14z^{(3+a)(3+b)}$

Proofs:

$$\begin{aligned}
 M_1(G, z) &= \sum_{uv \in E(G)} z^{[d_u + d_v]} \\
 &= |E_{\{1,3\}}(G)| z^4 + |E_{\{2,3\}}(G)| z^5 + |E_{\{3,3\}}(G)| z^6 \\
 &= 10z^4 + 22z^5 + 14z^6
 \end{aligned}$$

$$\begin{aligned}
 1. \quad M_2(G, z) &= \sum_{uv \in E(G)} z^{[d_u \times d_v]} \\
 &= |E_{\{1,3\}}(G)| z^3 + |E_{\{2,3\}}(G)| z^6 + |E_{\{3,3\}}(G)| z^9 \\
 &= 10z^3 + 22z^6 + 14z^9
 \end{aligned}$$

$$\begin{aligned}
 2. \quad M_3(G, z) &= \sum_{uv \in E(G)} z^{[d_u - d_v]} \\
 &= |E_{\{1,3\}}(G)| z^2 + |E_{\{2,3\}}(G)| z^1 + |E_{\{3,3\}}(G)| z^0 \\
 &= 10z^2 + 22z + 14
 \end{aligned}$$

$$\begin{aligned}
 3. \quad M_4(G, z) &= \sum_{uv \in E(G)} z^{d_u [d_u + d_v]} \\
 &= |E_{\{1,3\}}(G)| z^4 + |E_{\{2,3\}}(G)| z^{10} + |E_{\{3,3\}}(G)| z^{18} \\
 &= 10z^4 + 22z^{10} + 14z^{18}
 \end{aligned}$$

$$\begin{aligned}
 4. \quad M_5(G, z) &= \sum_{uv \in E(G)} z^{d_u + d_v} \\
 &= |E_{\{1,3\}}(G)| z^{12} + |E_{\{2,3\}}(G)| z^{15} + |E_{\{3,3\}}(G)| z^{18} \\
 &= 10z^{12} + 22z^{15} + 14z^{18} \\
 5. \quad M_{a,b}(G, z) &= \sum_{uv \in E(G)} z^{[ad_u + bd_v]} \\
 &= |E_{\{1,3\}}(G)| z^{a+3b} + |E_{\{2,3\}}(G)| z^{2a+3b} + |E_{\{3,3\}}(G)| z^{3a+3b} \\
 &= 10z^{a+3b} + 22z^{2a+3b} + 14z^{3a+3b} \\
 6. \quad M'_{a,b}(G, z) &= \sum_{uv \in E(G)} z^{(d_u + a)(b + d_v)} \\
 &= |E_{\{1,3\}}(G)| z^{(1+a)(3+b)} + |E_{\{2,3\}}(G)| z^{(2+a)(3+b)} + \\
 &\quad |E_{\{3,3\}}(G)| z^{(3+a)(3+b)} \\
 &= 10z^{(1+a)(3+b)} + 22z^{(2+a)(3+b)} + 14z^{(3+a)(3+b)}
 \end{aligned}$$

Theorem 2.12.

Let G be the graph of theaflavin. The redefined Zagreb indices are given below.

1. $ReZG_1 = 41$
2. $ReZG_2 = 40.9$
3. $ReZG_3 = 1536$

Proofs:

$$\begin{aligned}
 1. \quad ReZG_1 &= \sum_{uv \in E(G)} \frac{d_u + d_v}{d_u \times d_v} \\
 &= (10) \left(\frac{4}{3} \right) + (22) \left(\frac{5}{6} \right) + (14) \left(\frac{6}{9} \right) \\
 &= 41 \\
 2. \quad ReZG_2 &= \sum_{uv \in E(G)} \frac{d_u \times d_v}{d_u + d_v} \\
 &= (10) \left(\frac{3}{4} \right) + (22) \left(\frac{6}{5} \right) + (14) \left(\frac{9}{6} \right) \\
 &= 40.9 \\
 3. \quad ReZG_3 &= \sum_{uv \in E(G)} (d_u + d_v) (d_u \times d_v) \\
 &= (10) (3 \times 4) + (22) (6 \times 5) + (14) (9 \times 6) \\
 &= 1536
 \end{aligned}$$

3. Conclusion

In this article, we calculated the Zagreb polynomials and the redefined Zagreb indices for the chemical compounds used in the treatment of COVID-19. It has been demonstrated with certainty that the TIs help to anticipate their numerous properties without setting off to the wet lab.

References

- [1] Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020 Feb 15;395(10223):497-506. [https://doi.org/10.1016/S01406736\(20\)30183-5](https://doi.org/10.1016/S01406736(20)30183-5)
- [2] Wang M, Cao R, Zhang L, et al. Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. *Cell Res*. 2020 Mar;30(3):269-71.
- [3] Zhou D, Dai SM, Tong Q. COVID-19: a recommendation to examine the effect of hydroxychloroquine in preventing infection and progression. *J Antimicrob Chemother*. 2020 Jul 1;75(7):1667-70. <https://doi.org/10.1093/jac/dkaa114>
- [4] Lung J, Lin YS, Yang YH, et al. The potential chemical structure of anti-SARS-CoV-2 RNA-dependent RNA polymerase. *J Med Virol*. 2020 Jun;92(6):693-7. <https://doi.org/10.1002/jmv.25761>
- [5] Morse JS, Lalonde T, Xu S, et al. Learning from the past: possible urgent prevention and treatment options for severe acute respiratory infections caused by 2019-nCoV. *Chembiochem*. 2020 Mar 2;21(5):730-8. <https://doi.org/10.1002/cbic.202000047>
- [6] Xu X, Chen P, Wang J, et al. Evolution of the novel coronavirus from the ongoing Wuhan outbreak and modeling of its spike protein for risk of human transmission. *Sci China Life Sci*. 2020 Mar;63(3):457-60.
- [7] Warren TK, Jordan R, Lo MK, et al. Therapeutic efficacy of the small molecule GS-5734 against Ebola virus in rhesus monkeys. *Nature*. 2016 Mar;531(7594):381-5.
- [8] Gutman I, Ružić B, Trinajstić N, et al. Graph theory and molecular orbitals. XII. Acyclic polyenes. *J Chem Phys*. 1975 May 1;62(9):3399-405. <https://doi.org/10.1063/1.430994>

- [9] Fath-Tabar GH. Old and new Zagreb indices of graphs. *MATCH Commun Math Comput Chem*. 2011 Jan 1;65(1):79-84.
- [10] Ranjini PS, Lokesha V, Bindusree AR, et al. New bounds on Zagreb indices and the Zagreb co-indices. *Boletim da Sociedade Paranaense de Matemática*. 2013;31(1):51-5.
- [11] Fath-Tabar G. Zagreb polynomial and Pi Indices of some nano structures. *Dig J Nanomater Biostruct (DJNB)*. 2009 Mar 1;4(1):189-191.
- [12] Bindusree AR, Cangul IN, Lokesha V, Cevik AS. Zagreb polynomials of three graph operators. *Filomat*. 2016 Jan 1;30(7):1979-86.
- [13] Ranjini PS, Lokesha V, Usha A. Relation between phenylene and hexagonal squeezeusing harmonic index. *Int J Graph Theory*. 2013;1(4):116-21.
- [14] Gao W, Younas M, Farooq A, Virk AU, Nazeer W. Some reverse degree-based topological indices and polynomials of dendrimers. *Math*. 2018 Oct;6(10):214. <https://doi.org/10.3390/math6100214>
- [15] Gao W, Wang W, Dimitrov D, et al. Nano properties analysis via fourth multiplicative ABC indicator calculating. *Arabian J Chem*. 2018 Sep 1;11(6):793-801. <https://doi.org/10.1016/j.arabjc.2017.12.024>
- [16] Kang SM, Zahid MA, Nazeer W, et al. Calculating the degree-based topological indices of dendrimers. *Open Chem*. 2018 Aug 3;16(1):681-8.
- [17] Shao Z, Virk AU, Javed MS, et al. Degree based graph invariants for the molecular graph of Bismuth Tri-Iodide. *Engi Appl Sci Lett*. 2019 Dec;2(1):01-11.
- [18] Gao W, Wang W, Farahani MR. Topological indices study of molecular structure in anticancer drugs. *Journal of chemistry*. 2016 Feb;2016. <https://doi.org/10.1155/2016/3216327>
- [19] Virk AU. Some New Topological Invariants for Chemical Structures Used in the Treatment of COVID-19 Patients. *Turk J Math Comput Sci*.;12(2):112-9. <https://doi.org/10.47000/tjmcs.730730>